

NEWS2 - NB NEW ONSET CONFUSION SCORES 3 UNDER 'CONSCIOUSNESS'

Physiological parameter	Score						
	3	2	1	0	1	2	3
Respiration rate (per minute)	≤8		9–11	12–20		21–24	≥25
SpO ₂ Scale 1 (%)	≤91	92–93	94–95	≥96			
SpO ₂ Scale 2 (%)	≤83	84–85	86–87	88–92 ≥93 on air	93–94 on oxygen	95–96 on oxygen	≥97 on oxygen
Air or oxygen?		Oxygen		Air			
Systolic blood pressure (mmHg)	≤90	91–100	101–110	111–219			≥220
Pulse (per minute)	≤40		41–50	51–90	91–110	111–130	≥131
Consciousness				Alert			CVPU
Temperature (°C)	≤35.0		35.1–36.0	36.1–38.0	38.1–39.0	≥39.1	

ROCKWOOD CLINICAL FRAILTY SCALE

Clinical Frailty Scale*



1 Very Fit – People who are robust, active, energetic and motivated. These people commonly exercise regularly. They are among the fittest for their age.



2 Well – People who have **no active disease symptoms** but are less fit than category 1. Often, they exercise or are very **active occasionally**, e.g. seasonally.



3 Managing Well – People whose **medical problems are well controlled**, but are **not regularly active** beyond routine walking.



4 Vulnerable – While **not dependent** on others for daily help, often **symptoms limit activities**. A common complaint is being "slowed up", and/or being tired during the day.



5 Mildly Frail – These people often have **more evident slowing**, and need help in **high order IADLs** (finances, transportation, heavy housework, medications). Typically, mild frailty progressively impairs shopping and walking outside alone, meal preparation and housework.



6 Moderately Frail – People need help with **all outside activities** and with **keeping house**. Inside, they often have problems with stairs and need **help with bathing** and might need minimal assistance (cuing, standby) with dressing.



7 Severely Frail – **Completely dependent for personal care**, from whatever cause (physical or cognitive). Even so, they seem stable and not at high risk of dying (within ~ 6 months).



8 Very Severely Frail – Completely dependent, approaching the end of life. Typically, they could not recover even from a minor illness.



9. Terminally Ill - Approaching the end of life. This category applies to people with a **life expectancy <6 months**, who are **not otherwise evidently frail**.

Scoring frailty in people with dementia

The degree of frailty corresponds to the degree of dementia. Common **symptoms in mild dementia** include forgetting the details of a recent event, though still remembering the event itself, repeating the same question/story and social withdrawal.

In **moderate dementia**, recent memory is very impaired, even though they seemingly can remember their past life events well. They can do personal care with prompting.

In **severe dementia**, they cannot do personal care without help.

* 1. Canadian Study on Health & Aging, Revised 2008.
2. K. Rockwood et al. A global clinical measure of fitness and frailty in elderly people. CMAJ 2005;173:489-495.

Scoring charts for SAMBA23 Patient Data Collection

Links for calculating Index of Multiple Deprivation decile:

England: [English indices of deprivation 2019: Postcode Lookup \(opendatacommunities.org\)](https://www.opendatacommunities.org/england/indices-of-deprivation-2019/postcode-lookup)

Scotland: <https://simd.scot/#/simd2020/BTTTT/9/-4.0000/55.9000/>

Wales: <https://wimd.gov.wales>

Northern Ireland: <https://deprivation.nisra.gov.uk/>